



Genome and Plasmid Sequences of Escherichia coli KV7, an Extended-Spectrum -Lactamase Isolate Derived from Feces of a Healthy Pig

Bateman, Michael D; de Vries, Stefan P W; Gupta, Srishti; Guardabassi, Luca; Cavaco, Lina ; Grant, Andrew J; Holmes, Mark A

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.00595-17](https://doi.org/10.1128/genomeA.00595-17)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Bateman, M. D., de Vries, S. P. W., Gupta, S., Guardabassi, L., Cavaco, L., Grant, A. J., & Holmes, M. A. (2017). Genome and Plasmid Sequences of Escherichia coli KV7, an Extended-Spectrum -Lactamase Isolate Derived from Feces of a Healthy Pig. *Genome Announcements*, 5(38), [e00595-17].
<https://doi.org/10.1128/genomeA.00595-17>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Genome and Plasmid Sequences of *Escherichia coli* KV7, an Extended-Spectrum β -Lactamase Isolate Derived from Feces of a Healthy Pig

Michael D. Bateman,^a Stefan P. W. de Vries,^{a*} Srishti Gupta,^a Luca Guardabassi,^b
Lina M. Cavaco,^c Andrew J. Grant,^a Mark A. Holmes^a

Department of Veterinary Medicine, University of Cambridge, Cambridge, United Kingdom^a; Ross University School of Veterinary Medicine, Basseterre, St. Kitts, West Indies^b; National Food Institute, Technical University of Denmark, Kongens Lyngby, Denmark^c

ABSTRACT We present single-contig assemblies for *Escherichia coli* strain KV7 (serotype O27, phylogenetic group D) and its six plasmids, isolated from a healthy pig, as determined by PacBio RS II and Illumina MiSeq sequencing. The chromosome of 4,997,475 bp and G+C content of 50.75% harbored 4,540 protein-encoding genes.

A 2007 study (1) of healthy Danish pigs under prophylactic ceftiofur treatment uncovered a number of distinct strains of *Escherichia coli* carrying the *bla*_{CTX-M-1} gene and suggested that CTX-M-1 resistance spreads horizontally rather than clonally. A spontaneous nalidixic acid-resistant derivative of one of these strains that produces extended-spectrum β -lactamase (serotype 27, phylogenetic group D), named KV7, was investigated in a further study (2), showing that such strains thrive in pigs under treatment with several β -lactams. Sequencing of this isolate was performed in order to provide more detailed genetic data of a strain used for experimental infections of pigs.

The genomic and plasmid DNA was isolated from early exponential-phase cultures (LB medium) with Genomic-tip 100/G columns (Qiagen) and the plasmid midikit (Qiagen), respectively. Sequencing was performed using Illumina MiSeq and Pacific Biosciences technologies. Prior to PacBio RS II sequencing, the DNA was cleaned with a phenol extraction, and each preparation was sequenced on a single single-molecule real-time (SMRT) cell (Earlham Institute, Norwich, United Kingdom). For Illumina sequencing, DNA was sheared in an M220-focused ultrasonicator (Covaris). Sequencing libraries were prepared using the NEBNext Ultra II DNA library prep kit (New England Biolabs). Sequencing was performed using 250-bp paired-end sequencing on the MiSeq platform (V3 chemistry).

Assembly of genomic and plasmid DNA PacBio sequence reads was carried out at the Earlham Institute using the Hierarchical Genome Assembly Process (HGAP.3). Combining results from PacBio assemblies yielded single contigs for the chromosome and for five plasmids. A sixth plasmid was identified manually using a combination of raw reads and assembly fragments from both PacBio and Illumina sequencing assemblies; assembly of the Illumina reads was done with plasmidSPAdes version 3.9.0 (3, 4). This plasmid (P6) harbored a large repetitive region spanning approximately 2 kb. Polishing at the single-nucleotide polymorphism level was performed using Illumina reads with Snippy version 3.1. We did not find evidence of further plasmids.

The strain has multilocus sequence types (MLST) ST57 and ST533 using the original and enhanced *E. coli* MLST schemes, respectively. The KV7 chromosome was 4,997,475 bp with a GC content of 50.75%. The plasmid sizes were 223,014 bp (P1), 111,092 bp (P2), 71,546 bp (P3), 56,562 bp (P4), 32,688 bp (P5), and 39,510 bp (P6). Automated annotation was obtained with Prokka version 1.11 at Galaxy Queensland

Received 8 May 2017 Accepted 10 August 2017 Published 21 September 2017

Citation Bateman MD, de Vries SPW, Gupta S, Guardabassi L, Cavaco LM, Grant AJ, Holmes MA. 2017. Genome and plasmid sequences of *Escherichia coli* KV7, an extended-spectrum β -lactamase isolate derived from feces of a healthy pig. *Genome Announc* 5:e00595-17. <https://doi.org/10.1128/genomeA.00595-17>.

Copyright © 2017 Bateman et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mark A. Holmes, mah1@cam.ac.uk.

* Present address: Stefan P. W. de Vries, Host-Microbe Interactomics, Department of Animal Sciences, Wageningen University and Research, Wageningen, the Netherlands.

M.D.B. and S.P.W.D.V. contributed equally to this work.

(5), revealing the presence of 4,540 chromosomal protein-encoding genes. In addition to *bla*_{CTX-M-1}, a variety of antimicrobial resistance genes were identified on both the chromosome and several of the plasmids using BLAST together with the ARG-ANNOT database (6). These included genes predicted to encode resistance to aminoglycosides (*strA*, *strB*, and *aadA*), β -lactams (*bla*_{CTX-M-1}, *bla*_{TEM-1D}, *ampC1*, *ampC2*, and *ampH* and genes encoding penicillin-binding proteins), chloramphenicol (*catA1* and *cmlA*), a macrolide (*mphA*), sulfonamides (*sulI* and *sulII*), tetracycline (*tetB*), and trimethoprim (*dfrA*).

Accession number(s). The complete genome sequences of *E. coli* KV7 and plasmids have been deposited in the European Nucleotide Archive (ENA) (<http://www.ebi.ac.uk/ena>) and GenBank under accession no. [LT795502](#) for the KV7 genome and [LT795503](#), [LT795504](#), [LT795505](#), [LT795506](#), [LT795507](#), and [LT795508](#) for the plasmids (study no. PRJEB19461).

ACKNOWLEDGMENT

This work was supported by Medical Research Council (UK) grants MR/N002660/1 and G1001787/1.

REFERENCES

1. Jørgensen CJ, Cavaco LM, Hasman H, Emborg HD, Guardabassi L. 2007. Occurrence of CTX-M-1-producing *Escherichia coli* in pigs treated with ceftiofur. *J Antimicrob Chemother* 59:1040–1042. <https://doi.org/10.1093/jac/dkm075>.
2. Cavaco LM, Abatih E, Aarestrup FM, Guardabassi L. 2008. Selection and persistence of CTX-M-producing *Escherichia coli* in the intestinal flora of pigs treated with amoxicillin, ceftiofur, or cefquinome. *Antimicrob Agents Chemother* 52:3612–3616. <https://doi.org/10.1128/AAC.00354-08>.
3. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
4. Antipov D, Hartwick N, Shen M, Raiko M, Lapidus A, Pevzner PA. 2016. plasmidSPAdes: assembling plasmids from whole genome sequencing data. *Bioinformatics* 32:3380–3387. <https://doi.org/10.1093/bioinformatics/btw493>.
5. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
6. Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, Rolain JM. 2014. ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrob Agents Chemother* 58:212–220. <https://doi.org/10.1128/AAC.01310-13>.